

6

Magnetic Field Effects on Free Radical Reactions in Biology

Stefan Engström

CONTENTS

6.1	Introduction	157
6.2	Theoretical Background	158
6.2.1	Hyperfine Interaction-Induced Singlet-to-Triplet Conversion.....	159
6.2.2	High-Field Regime: Spin Rephrasing through the Δg Mechanism	159
6.2.3	Low-Field Effect: $B < 1$ mT	160
6.2.4	Free Radicals in Radio Frequency Fields.....	160
6.3	General Characteristics of the Free Radical Mechanism.....	160
6.3.1	Experimental Discrimination of Free Radical Models.....	161
6.4	Free Radicals in Biology	161
6.4.1	Biological Transduction Mechanisms.....	161
6.4.2	Role of Freely Diffusing Radicals.....	162
6.4.3	Animal Navigation Models Based on Free Radicals	163
6.4.4	Coenzyme B ₁₂ -Dependent Reactions.....	163
6.4.5	Other Experimental Observations.....	163
6.5	Conclusion	164
	References	164

6.1 Introduction

The physical chemistry of spin-correlated free radical pairs offers several mechanisms explaining how magnetic fields may influence biochemical processes. The mechanisms are classified on the basis of the dominating contribution to spin interconversion, and they cover a wide range of field strengths. Of particular interest is what is called the low-field mechanism, which has been extensively developed over the last decade and is now capable of explaining biological effects induced by magnetic fields well below 1 mT.

The principal mechanism behind the free radical mechanism was discovered in the physical problem of magnetic field dependence on positronium decay (Deutsch and Brown, 1952; Halpern, 1954). However, the development of the radical pair mechanism in chemistry has its roots in the work of Kaptein and Oosterhoff (1969), Closs (1969), and Brocklehurst (1969).

An ambitious survey of the literature up to its date of publication is the review of Steiner and Ulrich (1989), which lists some 775 references, 58 of which are themselves reviews on magnetokinetic phenomena. Another, now classic, reference on the subject is the book by Salikhov et al. (1984). McLauchlan and Steiner (1991) published a review including the possible mechanisms at lower fields. The review by Grissom (1995) explored the higher-field mechanisms with particular attention to the context of biological systems. A didactic paper geared toward the issues in biological systems is that of Brocklehurst and McLauchlan (1996). There are also some recent reviews on the free radical mechanism in general (Woodward, 2002) and with particular attention to biological systems (Brocklehurst, 2002).

6.2 Theoretical Background

A radical is an atom or a molecule with an unpaired electron. It tends to be highly reactive, a property that defines their best known roles in biology. A spin-correlated (or geminate) radical pair is typically created by hemolytic cleavage of a covalent bond, that is, each molecule retains one of the electrons that formed the chemical bond that was broken. The electron spins may remain correlated for a significant time (microseconds) after the pair's creation. As the radicals separate, the electron interaction term becomes small, and the electron states of the pair will fluctuate between antiparallel (singlet, or "S") and parallel (triplet, or "T") because of coherent spin evolution by hyperfine interactions between the electron spin and the nuclei. There is a chance of reencounter between the spin-correlated radicals. Reforming the bond is only permitted by quantum spin selection rules if the electron spins are oriented in the singlet state.


Magnetic fields can affect the electron state in the time before the reencounter; and if singlet and triplet pairs have different chemical fates, we have a basis for magnetic field effects on free radical chemistry (Figure 6.1). In exemplifying the process, we will assume that the radical pair is formed in the singlet state, the normal case for biologically relevant reactions (Eveson et al., 2000).

Many physical transduction mechanisms proposed to explain magnetic field effects in biology are very vulnerable to the randomizing effects of thermal noise at normal temperatures of living systems (Binhi and Savin, 2003). The free radical mechanism is uniquely resistant to this obstacle since it is a nuclear effect and is not strongly coupled to the thermal bath (Adair, 1999).

A quantum mechanical formulation of the free radical mechanism contains many possible contributions to the hamiltonian; Steiner and Ulrich (1989) provide a good categorization of the main components. The stochastic Liouville equation (SLE) is a tool

FIGURE 6.1

A molecule "C" (a) is split into two radicals (b). After diffusion and spin interconversion (c), the radicals may reencounter while still spin correlated (d). If the encounter occurs in the singlet state (e, top), the radicals may recombine. If the encounter occurs in the triplet state (e, bottom), recombination cannot occur, and the radicals will diffuse apart again and eventually lose their spin correlation. A magnetic field can influence this reaction by changing the rate of spin interconversion as long as the singlet and triplet products have different chemical fates.



for addressing the problem of simultaneous spin mixing and diffusion, but simplified models in which these two components are treated separately are often useful for the great reduction in problem complexity (Brocklehurst and McLauchlan, 1996). Recently, analytical results using a backward SLE have been presented (Pedersen and Christensen, 2004).

6.2.1 Hyperfine Interaction-Induced Singlet-to-Triplet Conversion

Hyperfine interactions between the spins of the electron and the nucleus cause the electron spins of the radicals to precess and induce singlet-to-triplet conversion. The triplet state has a net magnetic moment, and in the presence of an external magnetic field the energy levels of the triplet states that have a moment aligned with the magnetic field (T_+ and T_-) will be separated by Zeeman splitting. As the applied field strength is increased, the T_+ and T_- energy levels will be shifted away from the singlet state so much that they are decoupled from the spin interconversion process between singlet and triplet states, and only the remaining triplet state (T_0), which has a magnetic moment that is oriented perpendicular to the field, is capable of participating in the spin conversion process. In this way the magnetic field can reduce the number of triplet states that can be converted into singlet states and subsequently reform the original chemical bond. This is the “normal” magnetic field effect on free radical chemistry. It becomes relevant for external fields larger than the effective field driving the hyperfine interaction mixing, typically 1–10 mT.

A way in which singlet-to-triplet conversion can be facilitated is when an applied field causes the T_- energy level to cross the nonmagnetic singlet level, which occurs when the Zeeman differential matches the electron exchange interaction energy (cf. Figure 6.2). This effect has been observed for fields as low as 6.6 mT (Werner et al., 1993), but it can in principle be observed for much lower fields if the radical pairs are fixed with an appropriate separation.

6.2.2 High-Field Regime: Spin Rephrasing through the Δg Mechanism

The product of the magnetic field and the Landé g -factor determines the precession of the unpaired electron spin, independent of the hyperfine contribution. If the two radicals have slightly different g -factors, this provides an additional source of spin conversion. Differences are usually quite small, so this mechanism typically becomes significant only for quite large fields, $B > 0.1$ –1 T.




FIGURE 6.2

Energy level diagram for singlet and triplet states of a radical pair. At high fields, T_+ and T_- become completely disconnected from the singlet state, and only the T_0 state is left for singlet–triplet interconversion. At a specific B -field, the Zeeman split causes the T_- state to closely match the singlet state (circled), increasing the likelihood of transitions between the two states.

6.2.3 Low-Field Effect: $B < 1 \text{ mT}$

At fields below the hyperfine interaction energy, it is still possible to see effects of external fields under certain circumstances. It was found by Brocklehurst (1976) that the selection rules of the hyperfine-induced spin mixing are more restrictive in zero field than when a field is applied (McLauchlan and Steiner, 1991). This becomes relevant, even for a very small field, as long as the coherence of the pair's state is maintained for long periods (100 ns to 1 μ s). A helpful vector model to visualize this effect, along with some illustrative numerical examples, is given by Till et al. (1998).

The low-field effect (LFE) can theoretically produce a large (40%) drop in the singlet yield if the conditions are optimal (Timmel et al., 1998), but in practice only smaller effects attributed to this mechanism have been reported in the experimental literature.

Since long coherence times are required, it is necessary to understand spin relaxation effects and under what conditions they may be sufficiently long. Anisotropic hyperfine interactions provide noncoherent spin relaxation in solution, and it appears that the relaxation is slower in the low-field situation than has been generally thought (Fedin et al., 2001, 2003). It is becoming clear that understanding the local environment is crucial for evaluating LFEs (Eveson et al., 2000).

An important way to extend free radical magnetic field effects into the low-field region is to extend the lifetime of the spin-correlated pair. Integrating the magnetic field's influence over a longer time increases its ability to influence the spin evolution. Spin relaxation is not the primary problem here; rather, it is the reencounter probability that needs to be enhanced. There are several ways to achieve this:

1. Increased viscosity to restrict diffusion (Krissinel et al., 1999; Christensen and Pedersen, 2003; Kitahama et al., 2004)
2. Oppositely charged radicals that will oppose a tendency to diffuse apart (Adair, 1999)
3. Physically restricted mobility of the radicals through confining them to a surface or having the reaction taking place inside micelles, which are nanometer-scale compartments that form and reform on microsecond timescales and are able to confine the radical pairs and increase the reencounter probability (Eveson et al., 2000).

6.2.4 Free Radicals in Radio Frequency Fields

For higher-frequency fields ($f > 1 \text{ MHz}$) the time modulation of the field starts to correspond to timescales present in the reaction dynamics, and a range of resonant interactions become available (Timmel and Hore, 1996). A detailed treatment of the coenzyme B_{12} system demonstrated resonant phenomena for relatively low-level radio frequency (RF) fields using a variety of mathematical techniques (Canfield et al., 1994, 1995).

6.3 General Characteristics of the Free Radical Mechanism

Free radical reactions are generally quite fast. There is evidence of picosecond reactions (Gilch et al., 1998; Musewald et al., 1999), but many known reactions occur over nanosecond timescales. If the free radical diffusion is constrained by micelles (Eveson et al.,

2000; Christensen and Pedersen, 2003) or by Coulomb attraction (Horiuchi et al., 2003), it may be possible to extend the radical pair lifetimes to hundreds of nanoseconds or even microseconds.

Since the free radical mechanism is practically instantaneous when compared to the timescale of time-varying magnetic fields in the extremely low-frequency regime (<300 Hz), one would expect that the observable output from a biological detection of the field should depend only on the time-averaged absolute field amplitude (Scaiano et al., 1995). However, if there is a downstream system that is able to decode the low-frequency signal, this statement is not necessarily true (Engstrom, 1997; Engstrom and Fitzsimmons, 1999).

6.3.1 Experimental Discrimination of Free Radical Models

The relative orientation of a static and a much smaller oscillating field will provide a discriminating test for a free radical-based model as long as the timescale of the applied field is long compared to the lifetime of the radical pair. In an isotropic system, such as a liquid suspension, described by this field situation, only the amplitude of the magnetic field is relevant. While the smaller parallel field adds linearly to the larger static field, the perpendicular component is effectively reduced by the calculation of vector length and will cause a much smaller variation in field amplitude (Engstrom, 1997).

For oriented systems in which the free radical chemistry steps have spatial preferences, it is possible to have angular dependence with respect to the angle of the applied field, $f(\theta)$. One can argue that some symmetry properties are very likely to be present in that kind of situation (Ritz et al., 2000). Polarity changes are not expected to be relevant, so $f(\theta + 180^\circ) = f(\theta)$. Furthermore, due to the isotropic distribution of nuclear spins in the initial state of the radical pair, one also expects that $f(180^\circ - \theta) = f(\theta)$.

If the free radical mechanism is active for low-frequency fields, one can also expect a response to RF fields in the same system, a property not expected by any other suggested physical transduction mechanisms for magnetic fields (Henbest et al., 2004). In a comparison between static fields in the range 0–2 mT, it was shown that a 300- μ T, 5-MHz RF field applied perpendicularly or in parallel with the static field induced a response that was dependent on the magnitude of the static field (Henbest et al., 2004). This is qualitatively, if not quantitatively, similar to the observation that the magnetic sense of migratory birds can be disrupted by a 470-nT, 7-MHz field when the field is applied at an angle with the geomagnetic field (Ritz et al., 2004).

Magnetic isotope effects are another possible discriminating character of free radical effects, and it is possible to differentiate this signature from pure mass effects (Brocklehurst, 1997).

6.4 Free Radicals in Biology

6.4.1 Biological Transduction Mechanisms

Direct detection methods for free radicals have been developed in chemistry only relatively recently (Woodward, 2002). In biological systems, almost all experimental evidence for free radical involvement remains indirect. Generating hypotheses based on the signatures of free radical systems as outlined above are necessary to link free radical chemistry

to magnetic field effects, but there is circumstantial evidence that free radical chemistry underlies some effects reported in the bioelectromagnetics literature.

It has been suggested that complex dynamical systems may have special sensitivity to magnetic field influences (Grundler et al., 1992; Walleczek, 1995). This idea has been elaborated in a series of theoretical models of oscillatory systems (Eichwald and Walleczek, 1996a,b; Kaiser, 1996). These models show that enzyme dynamics involving free radical chemistry may be frequency specific, although only for timescales comparable to or slower than the chemical kinetics of the system (Eichwald and Walleczek, 1997). Field amplitude can influence enzyme dynamics in some instances and this can be used to exert some control over enzyme systems (Eichwald and Walleczek, 1998).

The peroxidase–oxidase system has interesting, well-documented dynamical properties (Scheeline et al., 1997). Detailed modeling of this system has shown how a magnetic field-induced perturbation can affect its dynamical behavior (Eichwald and Walleczek, 1998; Moller and Olsen, 1999). The point of interaction in this system is suspected to be electron-transfer enzyme intermediates (Moller and Olsen, 2000; Moller et al., 2000).

Downstream effects from changes in chemistry must be taken into account to evaluate biological significance (Brocklehurst and McLauchlan, 1996). This can both facilitate detection (Walleczek, 1995) as well as introduce interventions that could block the biological significance of a physical detection event.

It may be relevant that other enzyme systems have been studied in detail without the specific intention of addressing free radicals as a possible mechanism. Examples include electric and magnetic field effects in ATPase (Blank and Soo, 2001b). Direct interactions with DNA have been suggested; and electron transfer reactions are proposed interaction targets (Blank and Goodman, 2000; Blank and Soo, 2001a), although not by mechanisms addressed here. Myosin phosphorylation is another enzyme system that has shown sensitivity to time-varying (Markov et al., 1993) and static magnetic fields (Markov and Pilla, 1994), as well as gradient-specific effects (Engstrom et al., 2002).

6.4.2 Role of Freely Diffusing Radicals

Free radicals observed in biology are most commonly oxygen or nitrogen based with an unpaired electron, leading to the terms reactive oxygen species (ROS) and reactive nitrogen species (RNS). A dominant role for these radicals is to act in immunological defense. They are secreted by macrophages and neutrophils and during attempts to kill bacteria, viruses, and tumors (Nathan, 1992). The highly reactive nature of the radicals also means that damage to normal cells is possible, and various defense mechanisms against this have evolved as well (Yu, 1994). This immunological weapon with checks and balances already suggests that there is a signaling system built around free radicals, but it seems that the ROS and RNS also have roles in intracellular cell signaling (Lander, 1997) as well as intercellular communication (Thannickal and Fanburg, 2000).

Consider a biochemical reaction producing a spin-correlated free radical pair in the singlet state. Depending on the specific mechanism at work, the ratio of singlet-to-triplet product at reencounter will be modified. This will increase or decrease the fraction of pairs that tend to recombine because of a reencounter finding the spins in singlet versus the triplet states. For the LFE, the triplet state is favored, and we would see an excess of escape product. The situation is the opposite for the “normal” field effect, in which the T_- and T_+ states are decoupled from the interconversion process, increasing the proportion of singlet reencounters leading to a larger amount of cage product and leaving fewer freely diffusing radicals. The Δg mechanism brings the T_- and T_+ states back into play and therefore again boosts the triplet-reencounter escape products.

Given the wide involvement of free radicals in signaling and biological function, it is clear that there is the potential of both subtle and not-so-subtle effects on biological systems if we are able to alter the production of free radicals and thereby change the dynamics of already ongoing processes. The conventional wisdom regarding the deleterious effects of magnetic field effects on free radical recombination has been that more escape product means more radical-induced damage. This may be an oversimplification since the direct effects on cage or escape products are typically fairly small, certainly not larger than tens of percent, implying that drastic biological effects must involve downstream responses that amplify this relatively slight modulation. The answer may lie in the signaling properties of free radicals.

6.4.3 Animal Navigation Models Based on Free Radicals

The free radical mechanism was the first mechanism suggested as an explanation of avian use of magnetic fields for navigation (Schulten et al., 1978; Schulten, 1982). This model has since undergone several iterations of refinement (Ritz et al., 2000; Cintolesi et al., 2003). One interesting aspect of this work is a connection between photosensitivity and magnetoreception (Ritz et al., 2002). Dependence on light is a well-known feature of the avian magnetoreceptor (Deutschlander et al., 1999; Wiltschko et al., 2004a,b), but it has also appeared in other behavioral studies of animal magnetic field sensitivity (Prato et al., 1997, 1998).

The Ritz-Schulten model (Ritz et al., 2000) has an appealing geometrical application. Being integrated into the bird's retina, the suggested compass would appear as a modulation overlay on the bird's field of view. The mechanism operates through the so-called low field effect (LFE), based on a single nuclear spin, and operates near the limit of the theoretical sensitivity, despite omitting degrading effects such as the presence of multiple nuclear spins, dipolar effects, and various spin relaxation process that will start to become relevant for the long radical pair lifetimes (>100 ns) considered in the model.

Cryptochromes provide one possible source of free radicals in a spatially ordered system (Ritz et al., 2002). A recent theoretical model for avian magnetoreception develops that idea by investigating a flavin-tryptophan radical pair with a high degree of homology to the cryptochromes (Cintolesi et al., 2003). This multinucleus model is realistic in that it still manages to provide sensitivity to fields in the geomagnetic field range. Interestingly, it does not operate through the LFE described above (the multinucleus approach appears to remove most signs of that mechanism), but rather it depends on immobilized radicals and assumes that the free radical pair may have a lifetime up to 5 μ s.

6.4.4 Coenzyme B₁₂-Dependent Reactions

Magnetic field effects in the coenzyme B₁₂ are well explored experimentally with matching theoretical predictions (Harkins and Grissom, 1994; Grissom and Natarajan, 1997; Taoka et al., 1997). While most work on this model system has been concerned with intermediate and higher-field mechanisms, there are also detailed theoretical investigations suggesting that weak (<100 μ T), relatively low-frequency (<100 kHz), might be able to affect this system (Canfield et al., 1994, 1995, 1996).

6.4.5 Other Experimental Observations

The addition of iron ions or exposure to a 7-mT static magnetic field (SMF) did not affect the survival of rat lymphocytes *in vitro* when performed in isolation, but combined

exposure led to a significant increase in cell death (Jajte et al., 2002). One possible explanation of this behavior is that the addition of iron ions enhanced levels of ROS and that the field exposure further promoted the creation of free radicals, leading to cell death by both apoptosis and necrosis. An experiment with a similar rationale used added FeCl₂ to stimulate ROS production, and a 930-MHz, 5-W/m² cell phone-generated field affected a biological marker for ROS production. It should be noted that the vacuum magnetic field associated with this exposure is quite low (approximately 0.14 μT) and the frequency is a relatively unexplored region for this mechanism.

Proliferation of chick fibroblasts was observed to be enhanced by a 100-Hz, 0.7-mT sinusoidal magnetic field (Katsir et al., 1998). In a follow-up study it was found that free radical scavengers (Katsir and Parola, 1998) suppressed this effect, suggesting that that free radicals may have a role in mediating the magnetic field effect on proliferation.

Genotoxic effects from intermediate static magnetic fields (250 mT) have been studied in *Escherichia coli* DNA, both *in vivo* and *in vitro* (Potenza et al., 2004). Free radical formation was stimulated, and the genetic damage was mapped as a function of exposure duration. *In vitro* experiments showed detectable genotoxic effects, but the *in vivo* assays did not, indicating that cellular protective responses may prevent damage in the intact system.

A reported effect on the oxidative burst in neutrophils by a 0.1-mT field was attributed to free radicals (Roy et al., 1995). In that study, the connection to free radicals lies in that the fluorescent probe used to study the neutrophil activity reacts specifically with free radical-derived oxidants that create the fluorescing compounds. Work in neutrophils in humans (Heine et al., 1999) using a much larger field (1.5 T) did not find any effects of magnetic fields on the respiratory burst of human neutrophils or on the production of radical species.

Phagocytosis was observed to be affected by 0.5–1.5-mT, 50-Hz sinusoidal magnetic fields (Simko et al., 2001). An attendant increase in superoxide production may be an indication that the field stimulated the system through a free radical process.

6.5 Conclusion

Free radical reactions are ubiquitous in biology, and recent developments of the low-field mechanisms (Timmel et al., 2001) and the consideration of detailed biochemical systems (Cintolesi et al., 2003) make this mechanism a contender for field effects down to geo-magnetic field strengths. The physical transduction step is not vulnerable to thermal perturbations, a significant advantage over competing models.

This model does not produce large (factors >2) changes at the initial field detection step. Theoretical models and direct experimental observations in the low-field region typically operate around or below the 10% level, so we should expect the physical detection mechanism to need cooperation from downstream processes for biologically relevant detection of magnetic fields with free radicals as the starting point.

References

- Adair RK (1999): Effects of very weak magnetic fields on radical pair reformation. *Bioelectromagnetics* 20:255–263.
- Binh VN, Savin AV (2003): Effects of weak magnetic fields on biological systems: physical aspects. *Physics-Usppekhi* 46:259–291.

- Blank M, Goodman R (2000): Stimulation of the stress response by low-frequency electromagnetic fields: possibility of direct interaction with DNA. *IEEE Transactions on Plasma Science* 28:168–172.
- Blank M, Soo L (2001a): Electromagnetic acceleration of electron transfer reactions. *Journal of Cellular Biochemistry* 81:278–283.
- Blank M, Soo L (2001b): Optimal frequencies for magnetic acceleration of cytochrome oxidase and Na,K-ATPase reactions. *Bioelectrochemistry* 53:171–174.
- Brocklehurst B (1969): Formation of excited states by recombining organic ions. *Nature* 221:921.
- Brocklehurst B (1976): Spin correlation in geminate recombination of radical ions in hydrocarbons. 1. Theory of magnetic-field effect. *Journal of the Chemical Society—Faraday Transactions II* 72:1869–1884.
- Brocklehurst B (1997): Magnetic isotope effects in biology: a marker for radical pair reactions and electromagnetic field effects? *International Journal of Radiation Biology* 72:587–596.
- Brocklehurst B (2002): Magnetic fields and radical reactions: recent developments and their role in nature. *Chemical Society Reviews* 31:301–311.
- Brocklehurst B, McLauchlan KA (1996): Free radical mechanism for the effects of environmental electromagnetic fields on biological systems. *International Journal of Radiation Biology* 69:3–24.
- Canfield JM, Belford RL, Debrunner PG, Schulten KJ (1994): A perturbation-theory treatment of oscillating magnetic-fields in the radical pair mechanism. *Chemical Physics* 182:1–18 (see also *Journal of Chemical Physics* 191:347, 1995).
- Canfield JM, Belford RL, Debrunner PG, Schulten KJ (1995): A perturbation treatment of oscillating magnetic-fields in the radical pair mechanism using the Liouville equation. *Chemical Physics* 195:59–69.
- Canfield JM, Belford RL, Debrunner PG (1996): Calculations of Earth-strength steady and oscillating magnetic field effects in coenzyme B-12 radical pair systems. *Molecular Physics* 89:889–930.
- Christensen M, Pedersen JB (2003): On the validity of the one-particle diffusion model of geminate recombination in micelles. *Chemical Physics* 295:235–241.
- Cintolesi F, Ritz T, Kay CWM, Timmel CR, Hore PJ (2003): Anisotropic recombination of an immobilized photoinduced radical pair in a 50- μ T magnetic field: a model avian photomagnetoceptor. *Chemical Physics* 294:385–399.
- Closs GL (1969): A mechanism explaining nuclear spin polarizations in radical combination reactions. *Journal of the American Chemical Society* 91:4552.
- Deutsch M, Brown SC (1952): Zeeman effect and hyperfine splitting of positronium. *Physical Review* 85:1047–1048.
- Deutschlander ME, Phillips JB, Borland SC (1999): The case for light-dependent magnetic orientation in animals. *Journal of Experimental Biology* 202:891–908.
- Eichwald C, Walleczek J (1996a): Activation-dependent and biphasic electromagnetic field effects: model based on cooperative enzyme kinetics in cellular signaling. *Bioelectromagnetics* 17:427–435.
- Eichwald C, Walleczek J (1996b): Model for magnetic field effects on radical pair recombination in enzyme kinetics. *Biophysical Journal* 71:623–631.
- Eichwald C, Walleczek J (1997): Low-frequency-dependent effects of oscillating magnetic fields on radical pair recombination in enzyme kinetics. *Journal of Chemical Physics* 107:4943–4950.
- Eichwald C, Walleczek J (1998): Magnetic field perturbations as a tool for controlling enzyme-regulated and oscillatory biochemical reactions. *Biophysical Chemistry* 74:209–224.
- Engstrom S (1997): What is the time scale of magnetic field interaction in biological systems? *Bioelectromagnetics* 18:244–249.
- Engstrom S, Fitzsimmons R (1999): Five hypotheses to examine the nature of magnetic field transduction in biological systems. *Bioelectromagnetics* 20:423–430.
- Engstrom S, Markov MS, McLean MJ, Holcomb RR, Markov JM (2002): Effects of non-uniform static magnetic fields on the rate of myosin phosphorylation. *Bioelectromagnetics* 23:475–479.
- Eveson RW, Timmel CR, Brocklehurst B, Hore PJ, McLauchlan KA (2000): The effects of weak magnetic fields on radical recombination reactions in micelles. *International Journal of Radiation Biology* 76:1509–1522.

- Fedin MV, Purtov PA, Bagryanskaya EG (2001): Anisotropic hyperfine interaction-induced spin relaxation in a low magnetic field. *Chemical Physics Letters* 339:395–404.
- Fedin MV, Purtov PA, Bagryanskaya EG (2003): Spin relaxation of radicals in low and zero magnetic field. *Journal of Chemical Physics* 118:192–201.
- Gilch P, Pollinger-Dammer F, Musewald C, Michel-Beyerle ME, Steiner UE (1998): Magnetic field effect on picosecond electron transfer. *Science* 281:982–984.
- Grissom CB (1995): Magnetic-field effects in biology—a survey of possible mechanisms with emphasis on radical-pair recombination. *Chemical Reviews* 95:3–24.
- Grissom CB, Natarajan E (1997): Use of magnetic field effects to study coenzyme B-12-dependent reactions. *Vitamins and Coenzymes*, Pt K 281:235–247.
- Grundler W, Kaiser F, Keilmann F, Walleczek J (1992): Mechanisms of electromagnetic-interaction with cellular-systems. *Naturwissenschaften* 79:551–559.
- Halpern O (1954): Magnetic quenching of the positronium decay. *Physical Review* 94:904–907.
- Harkins TT, Grissom CB (1994): Magnetic-field effects on B-12 ethanolamine ammonia-lyase—evidence for a radical mechanism. *Science* 263:958–960.
- Heine J, Scheinichen D, Jaeger K, Herzog T, Sumpelmann R, Leuwer M (1999): Effect of magnetic resonance imaging on human respiratory burst of neutrophils. *FEBS Letters* 446:15–17.
- Henbest KB, Kukura P, Rodgers CT, Hore PJ, Timmel CR (2004): Radio frequency magnetic field effects on a radical recombination reaction: a diagnostic test for the radical pair mechanism. *Journal of the American Chemical Society* 126:8102–8103.
- Horiuchi M, Maeda K, Arai T (2003): Magnetic field effect on electron transfer reactions of flavin derivatives associated with micelles. *Applied Magnetic Resonance* 23:309–318.
- Jajte J, Grzegorzczak J, Zmyslony M, Rajkowska E (2002): Effect of 7 mT static magnetic field and iron ions on rat lymphocytes: apoptosis, necrosis and free radical processes. *Bioelectrochemistry* 57:107–111.
- Kaiser F (1996): External signals and internal oscillation dynamics: biophysical aspects and modeling approaches for interactions of weak electromagnetic fields at the cellular level. *Bioelectrochemistry and Bioenergetics* 41:3–18.
- Kaptein R, Oosterhoff LJ (1969): Chemically induced dynamic nuclear polarization III (anomalous multiplets of radical coupling and disproportionation products). *Chemical Physics Letters* 4:214–216.
- Katsir G, Parola AH (1998): Enhanced proliferation caused by a low frequency weak magnetic field in chick embryo fibroblasts is suppressed by radical scavengers. *Biochemical and Biophysical Research Communications* 252:753–756.
- Katsir G, Baram SC, Parola AH (1998): Effect of sinusoidally varying magnetic fields on cell proliferation and adenosine deaminase specific activity. *Bioelectromagnetics* 19:46–52.
- Kitahama Y, Wakasa M, Sakaguchi Y (2004): Viscosity dependence of the magnetic field effect due to the delta g mechanism. *Journal of Physical Chemistry A* 108:754–757.
- Krissinel EB, Burshtein AI, Lukzen NN, Steiner UE (1999): Magnetic field effect as a probe of distance-dependent electron transfer in systems undergoing free diffusion. *Molecular Physics* 96:1083–1097.
- Lander HM (1997): An essential role for free radicals and derived species in signal transduction. *FASEB Journal* 11:118–124.
- Markov MS, Pilla AA (1994): Static magnetic-field modulation of myosin phosphorylation—calcium-dependence in 2 enzyme preparations. *Bioelectrochemistry and Bioenergetics* 35(1–2):57–61.
- Markov MS, Wang S, Pilla AA (1993): Effects of weak low frequency sinusoidal and DC magnetic fields on myosin phosphorylation in a cell-free preparation. *Bioelectrochemistry and Bioenergetics* 30:119–125.
- McLauchlan KA, Steiner UE (1991): The spin-correlated radical pair as a reaction intermediate. *Molecular Physics* 73:241–263.
- Moller AC, Olsen LF (1999): Effect of magnetic fields on an oscillating enzyme reaction. *Journal of the American Chemical Society* 121:6351–6354.
- Moller AC, Olsen LF (2000): Perturbations of simple oscillations and complex dynamics in the peroxidase–oxidase reaction using magnetic fields. *Journal of Physical Chemistry B* 104:140–146.

- Moller AC, Lunding A, Olsen LF (2000): Further studies of the effect of magnetic fields on the oscillating peroxidase–oxidase reaction. *Physical Chemistry Chemical Physics* 2:3443–3446.
- Musewald C, Gilch P, Hartwich G, Pollinger-Dammer F, Scheer H, Michel-Beyerle ME (1999): Magnetic field dependence of ultrafast intersystem-crossing: a triplet mechanism on the picosecond time scale? *Journal of the American Chemical Society* 121:8876–8881.
- Nathan C (1992): Nitric-oxide as a secretory product of mammalian-cells. *FASEB Journal* 6:3051–3064.
- Pedersen JB, Christensen M (2004): The backward stochastic Liouville equation. *Journal of Physical Chemistry B* 108:9516–9523.
- Potenza L, Cucchiari L, Piatti E, Angelini U, Dacha M (2004): Effects of high static magnetic field exposure on different DNAs. *Bioelectromagnetics* 25:352–355.
- Prato FS, Kavaliers M, Cullen AP, Thomas AW (1997): Light-dependent and -independent behavioural effects of extremely low frequency magnetic fields in a land snail are consistent with a parametric resonance mechanism. *Bioelectromagnetics* 18:284–291.
- Prato FS, Kavaliers M, Thomas AW, Ossenkopp KP (1998): Modulatory actions of light on the behavioural responses to magnetic fields by land snails probably occur at the magnetic field detection stage. *Proceedings of the Royal Society of London Series B: Biological Sciences* 265:367–373.
- Ritz T, Adem S, Schulten K (2000): A model for photoreceptor-based magnetoreception in birds. *Biophysical Journal* 78:707–718.
- Ritz T, Dommer DH, Phillips JB (2002): Shedding light on vertebrate magnetoreception. *Neuron* 34:503–506.
- Ritz T, Thalau P, Phillips JB, Wiltschko R, Wiltschko W (2004): Resonance effects indicate a radical-pair mechanism for avian magnetic compass. *Nature* 429:177–180.
- Roy S, Noda Y, Eckert V, Traber MG, Mori A, Liburdy R, Packer L (1995): The phorbol 12-myristate 13-acetate (PMA)-induced oxidative burst in rat peritoneal neutrophils is increased by a 0.1 mT (60 Hz) magnetic-field. *FEBS Letters* 376:164–166.
- Salikhov KM, Molin YuN, Sagdeev RZ, Buchachenko AL (1984): *Spin Polarization and Magnetic Effects in Radical Reactions*. Elsevier, Amsterdam.
- Scaiano JC, Cozens FL, Mohtat N (1995): Influence of combined AC-DC magnetic-fields on free-radicals in organized and biological-systems—development of a model and application of the radical pair mechanism to radicals in micelles. *Photochemistry and Photobiology* 62:818–829.
- Scheeline A, Olson DL, Williksen EP, Horras GA, Klein ML, Larter R (1997): The peroxidase–oxidase oscillator and its constituent chemistries. *Chemical Reviews* 97:739–756.
- Schulten K (1982): Magnetic field effects in chemistry and biology. In Treusch J, ed. *Festkörperprobleme*, Vol. 22. Vieweg, Braunschweig, pp. 61–83.
- Schulten K, Swenberg CE, Weller A (1978): Biomagnetic sensory mechanism based on magnetic-field modulated coherent electron-spin motion. *Zeitschrift für Physikalische Chemie-Frankfurt* 111:1–5.
- Simko M, Droste S, Kriehuber R, Weiss DG (2001): Stimulation of phagocytosis and free radical production in murine macrophages by 50 Hz electromagnetic fields. *European Journal of Cell Biology* 80:562–566.
- Steiner UE, Ulrich T (1989): Magnetic-field effects in chemical-kinetics and related phenomena. *Chemical Reviews* 89:51–147.
- Taoka S, Padmakumar R, Grissom CB, Banerjee R (1997): Magnetic field effects on coenzyme B-12-dependent enzymes: validation of ethanolamine ammonia lyase results and extension to human methylmalonyl CoA mutase. *Bioelectromagnetics* 18:506–513.
- Thannickal VJ, Fanburg BL (2000): Reactive oxygen species in cell signaling. *American Journal of Physiology—Lung Cellular and Molecular Physiology* 279:L1005–L1028.
- Till U, Timmel CR, Brocklehurst B, Hore PJ (1998): The influence of very small magnetic fields on radical recombination reactions in the limit of slow recombination. *Chemical Physics Letters* 298:7–14.
- Timmel CR, Hore PJ (1996): Oscillating magnetic field effects on the yields of radical pair reactions. *Chemical Physics Letters* 257:401–408.
- Timmel CR, Till U, Brocklehurst B, McLauchlan KA, Hore PJ (1998): Effects of weak magnetic fields on free radical recombination reactions. *Molecular Physics* 95:71–89.
- Timmel CR, Cintolesi F, Brocklehurst B, Hore PJ (2001): Model calculations of magnetic field effects on the recombination reactions of radicals with anisotropic hyperfine interactions. *Chemical Physics Letters* 334:387–395.

- Walleczek J (1995): Magnetokinetic effects on radical pairs: a paradigm for magnetic field interactions with biological systems at lower than thermal energy. *Electromagnetic Fields* 250:395–420.
- Werner U, Kuhnle W, Staerk H (1993): Magnetic-field dependent reaction yields from radical-ion pairs linked by a partially rigid aliphatic chain. *Journal of Physical Chemistry* 97:9280–9287.
- Wiltschko W, Gesson M, Stapput K, Wiltschko R (2004a): Light-dependent magnetoreception in birds: interaction of at least two different receptors. *Naturwissenschaften* 91:130–134.
- Wiltschko W, Moller A, Gesson M, Noll C, Wiltschko R (2004b): Light-dependent magnetoreception in birds: analysis of the behaviour under red light after pre-exposure to red light. *Journal of Experimental Biology* 207:1193–1202.
- Woodward JR (2002): Radical pairs in solution. *Progress in Reaction Kinetics and Mechanism* 27:165–207.
- Yu BP (1994): Cellular defenses against damage from reactive oxygen species. *Physiological Reviews* 74:139–162.